

# Patients on GLP-1 medications\*

## STOP ALL

### Solid Foods at 5PM

When: 5PM the night before your procedure.

What to avoid: All solid foods, including snacks, gum, and candy.



## CLEAR LIQUIDS ONLY

Allowed from 5PM the day prior until 2 hours before your arrival time. Clear liquids include:

Water

Black tea & coffee (NO MILK/CREAM)

Fruit juice without pulp (like apple juice)

Gatorade or similar clear sports drinks



Stop drinking all clear liquids exactly 2 hours before your scheduled arrival time!


**Important – Failure to follow these instructions can lead to the cancellation of your procedure!**

# Pacientes que toman medicamentos GLP-1\*

**PARAR DE CONSUMIR TODOS LOS ALIMENTOS SÓLIDOS A LAS 5PM**



**Cuándo:** A las 5PM, la noche anterior a su procedimiento. **Qué evitar:** Todos los alimentos sólidos, incluyendo aperitivos, chicles y caramelos.



## **SOLO LÍQUIDOS CLAROS**

Permitido desde las 17:00 del día anterior hasta 2 horas antes de la hora de llegada.

Los líquidos claros incluyen:

Agua

Té y Café negro (SIN LECHE NI CREMA)

Jugo de frutas sin pulpa (por ejemplo, jugo de manzana)

Gatorade o bebidas deportivas claras similares



**Deje de beber todos los líquidos claros exactamente 2 horas antes de la hora programada de llegada.**

**Importante: No seguir estas instrucciones puede llevar a la cancelación de su procedimiento.**

# \*IMPORTANT!\*

Patients using WEEKLY Injectable GLP-1 Agonist Medications

Need to **STOP** at least ONE FULL WEEK before Surgery!

Examples Include:


- Semaglutide (Ozempic, Wegovy)
- Tirzepatide (Mounjaro, Zepbound)
  - Exanatide ER (Bydureon)
  - Albiglutide (Tanzeum)
  - Dulaglutide (Trulicity)

Patients on DAILY Injectable and Oral GLP-1 Agonist Medications

Need to **STOP** taking the day of Surgery!

Examples Include:

- Lixisenatide (Adlyxin)
- Semaglutide (Rybelsus, Wegovy)
  - Tirzepatide
- Orforglipron (Foundayo)
- Liraglutide (Victoza, Saxenda)
  - Exenatide (Byetta)



# ¡IMPORTANTE!

Pacientes que usan medicamentos GLP-1 Agonistas inyectables  
SEMANALES

¡Necesito PARAR al menos UNA SEMANA COMPLETA antes  
de la cirugía!

Ejemplos incluyen:

- Semaglutide (Ozempic, Wegovy)
- Tirzepatide (Mounjaro, Zepbound)
  - Exanatide ER (Bydureon)
  - Albiglutide (Tanzeum)
  - Dulaglutide (Trulicity)

Pacientes que toman DIARIAMENTE medicamentos inyectables y  
orales para agonistas GLP-1

¡Necesito DEJAR de tomar el día de la cirugía!

Ejemplos incluyen:

- Lixisenatide (Adlyxin)
- Semaglutide (Rybelsus, Wegovy)
  - Tirzepatide
- Orforglipron (Foundayo)
- Liraglutide (Victoza, Saxenda)
  - Exenatide (Byetta)

